

LA-3422

LA-3422

Bauer
D3

Reproduced From
Best Available Copy

A Facsimile Report



DISTRIBUTION STATEMENT A
Approved for Public Release
Distribution Unlimited

Reproduced by
UNITED STATES
ATOMIC ENERGY COMMISSION
Division of Technical Information
P.O. Box 62 Oak Ridge, Tennessee 37831

JUN 10 1966

Lovelace Foundation - Document Library
17822

20000915 039

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

This report expresses the opinions of the author or authors and does not necessarily reflect the opinions or views of the Los Alamos Scientific Laboratory.

BLANK PAGE

Printed in USA. Price ~~\$2.00~~. Available from the Clearinghouse for Federal Scientific and Technical Information, National Bureau of Standards, United States Department of Commerce, Springfield, Virginia

LA-3422
UC-34, PHYSICS
TID-4500 (46th Ed.)

LOS ALAMOS SCIENTIFIC LABORATORY
of the
University of California
LOS ALAMOS • NEW MEXICO

Report written: September 10, 1965

Report distributed: January 11, 1966

**Three-Dimensional Cartesian
Particle-in-Cell Calculations**

by

William R. Gage
Charles L. Mader

RELEASED FOR ARCHIVING
IN NUCLEAR SCIENCE ABSTRACTS

-1-

DISTRIBUTION STATEMENT A
Approved for Public Release
Distribution Unlimited

ABSTRACT

A study of the Particle-In-Cell method of numerically solving hydrodynamic problems in three dimensions is described. The closure of a cubical void in a cubical box was successfully computed. Various methods, including stereoscopic views, of pictorial description of the particle positions are described.

ACKNOWLEDGMENT

The authors gratefully acknowledge the assistance and contributions of F. H. Harlow of the Los Alamos Scientific Laboratory.

-3-

CONTENTS

Abstract	Page
Acknowledgment	3
I. Introduction	3
II. Computational Method	7
III. Void Closure	7
IV. Conclusions	8
Appendix A. The Hydrodynamic Equations	8
Appendix B. Stereo Plotting	37
Literature Cited	57
	62

FIGURES

1. A cube of nitromethane containing a centered cubical void.	9
2. The particle positions at various times for a cube of nitromethane, 0.024 cm on a side, containing a cubical void 0.008 cm on a side. The 0.024-cm side in the Z direction of the cube is cut into 12 equally spaced layers. A piston is applied to the bottom of the cube. Each group of 12 pictures presents views of the layers in succession as seen looking through the x,y face of the cube.	10

BLANK PAGE

3. The particle positions at various times for a cube of nitromethane containing a cubical void cut into 15 equally spaced layers in the Z direction. Each group of 15 pictures presents views of the successive layers through the void. 20
4. The particle positions through half of the void shown at various times in stereo projections looking down into the void at a 15° angle. 29
5. The particle positions through all of the void shown at various times in stereo projections looking down into the void at a 15° angle. 33

I. INTRODUCTION

The Particle-In-Cell method¹ for numerically solving two-dimensional hydrodynamic problems with chemical reaction and realistic equations of state has been remarkably successful in describing the formation of hot spots when shocks interact with density discontinuities^{2,3}. The Particle-In-Cell method allows large distortions to occur in the fluid and permits void closure. Because of these unique characteristics, the Particle-In-Cell method is the obvious one to use to solve the three-dimensional problem of shocks interacting with many voids. While sufficiently fast computers to resolve such a problem in detail will not be available for several years, we have investigated the Particle-In-Cell method in three dimensions to determine if the techniques used in the two-dimensional EIC code^{2,3} were adequate when extended to three dimensions. The results presented in this report show that the Particle-In-Cell method is adequate.

II. COMPUTATIONAL METHOD

The calculational technique used is the PTC (Particle-In-Cell) scheme described in detail in Appendix A.

The equation of state used to describe the fluid is the HON equation of state for nitromethane described in reference 2.

The calculations were performed using an IBM 7030 (STRETCH) code written as part of this study.

Accuracy in resolving the dynamics is dependent upon the number of cells and particles. For practical purposes, the accuracy is limited by the amount of machine time that is available. In the calculations reported here, 1728 cells were used with 12 cells in each of the X, Y, Z directions. We used 16 particles per cell, or a total of 27,648 particles. Such a crude mesh is adequate for studying the method, but would not give sufficient resolution for most problems of interest.

III. VOID CLOSURE

The model studied was a cube of nitromethane with 12 cells on a side containing a cubical void with 4 cells on a side. The model is shown in Figure 1 with the coordinate system which was used. A piston is applied to the bottom of the cube, shocking the nitromethane to 85 kbar.

We also studied the reactive flow problem of a 125-kbar shock driving a detonation through the cube of nitromethane. The results compared well with those obtained in one and two dimensions.

The particle positions of each of the 12 layers of cells in the Z direction are shown in Figure 2.

The particle positions of each of 16 equally spaced layers through the void in the Z direction are shown in Figure 3.

Stereo projections of half the cells and all the cells through the void are shown in Figures 4 and 5. With a viewer, a stereo effect is achieved. The angle of observation and particular amount of information to be displayed depends upon the problem; however, stereo projections are a useful method of presentation of the results, and the method is described in Appendix B.

Perhaps the most useful method of presenting the results is to make clear prints of the 24 layers of particles in half of the cube and mount them to scale in grooved Lucite boxes. By this technique, one obtains a three-dimensional scale model.

IV. CONCLUSIONS

The Particle-In-Cell method has been successfully extended to three dimensions. Various methods of pictorial description of the results were tried. Each method had its uses, with the three-dimensional model being the most informative, but least convenient. Solution of three-dimensional hydrodynamic problems of practical interest appears to be possible as soon as computers with sufficient speed (at least 100 times faster than STRETCH) are available.








Fig. 1. A cube of nitromethane containing a centered cubical void.


Figure 2

The particle positions at various times for a cube of nitromethane, 0.024 cm on a side, containing a cubical void 0.008 cm on a side. The 0.024-cm side in the Z direction of the cube is cut into 12 equally spaced layers. A piston is applied to the bottom of the cube. Each group of 12 pictures presents views of the layers in succession as seen looking through the x,y face of the cube.







1
 2
 3
 4
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525




1




.....





-15-




1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2620, 2621, 2622, 2623, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2638, 2639, 2640, 2641, 2642, 2643, 2644, 2645, 2646, 2647, 2648, 2649, 2650, 2651, 2652, 2653, 2654, 2655, 2656, 2657, 2658, 2659, 2660, 2661, 2662, 2663, 2664, 2665, 2666, 2667, 2668, 2669, 2670, 2671, 2672, 2673, 2674, 2675, 2676, 2677, 2678, 26




[illegible][illegible]


-74-




THE LEFT HALF OF THE IMAGE IS A DENSE, DIAGONAL HATCHING PATTERN. THE RIGHT HALF IS A SOLID BLACK BACKGROUND.




THE LEFT HALF OF THE IMAGE IS A DENSE, DIAGONAL HATCHING PATTERN. THE RIGHT HALF IS A SOLID BLACK BACKGROUND.




THE LEFT HALF OF THE IMAGE IS A DENSE, DIAGONAL HATCHING PATTERN. THE RIGHT HALF IS A SOLID BLACK BACKGROUND.




THE LEFT HALF OF THE IMAGE IS A DENSE, DIAGONAL HATCHING PATTERN. THE RIGHT HALF IS A SOLID BLACK BACKGROUND.




THE LEFT HALF OF THE IMAGE IS A DENSE, DIAGONAL HATCHING PATTERN. THE RIGHT HALF IS A SOLID BLACK BACKGROUND.




THE LEFT HALF OF THE IMAGE IS A DENSE, DIAGONAL HATCHING PATTERN. THE RIGHT HALF IS A SOLID BLACK BACKGROUND.




THE LEFT HALF OF THE IMAGE IS A DENSE, DIAGONAL HATCHING PATTERN. THE RIGHT HALF IS A SOLID BLACK BACKGROUND.




THE LEFT HALF OF THE IMAGE IS A DENSE, DIAGONAL HATCHING PATTERN. THE RIGHT HALF IS A SOLID BLACK BACKGROUND.




THE LEFT IS A REPRESENTATION OF THE RIGHT SIDE OF THE LEFT SIDE




THE LEFT IS A REPRESENTATION OF THE RIGHT SIDE OF THE LEFT SIDE




THE LEFT IS A REPRESENTATION OF THE RIGHT SIDE OF THE LEFT SIDE




THE LEFT IS A REPRESENTATION OF THE RIGHT SIDE OF THE LEFT SIDE




THE LEFT IS A REPRESENTATION OF THE RIGHT SIDE OF THE LEFT SIDE



THE LEFT IS A REPRESENTATION OF THE RIGHT SIDE OF THE LEFT SIDE






THE LEFT IS A REPRESENTATION OF THE RIGHT SIDE OF THE LEFT SIDE




THE LEFT IS A REPRESENTATION OF THE RIGHT SIDE OF THE LEFT SIDE

The particle positions at various times for a cube of nitromethane containing a cubical void cut into 16 equally spaced layers in the Z direction. Each group of 16 pictures presents views of the successive layers through the void.







THESE ARE THE RESULTS OF THE TESTS CONDUCTED ON THE 11/11/1994




THESE ARE THE RESULTS OF THE TESTS CONDUCTED ON THE 11/11/1994




THESE ARE THE RESULTS OF THE TESTS CONDUCTED ON THE 11/11/1994




THESE ARE THE RESULTS OF THE TESTS CONDUCTED ON THE 11/11/1994




THESE ARE THE RESULTS OF THE TESTS CONDUCTED ON THE 11/11/1994



THESE ARE THE RESULTS OF THE TESTS CONDUCTED ON THE 11/11/1994



THESE ARE THE RESULTS OF THE TESTS CONDUCTED ON THE 11/11/1994



THESE ARE THE RESULTS OF THE TESTS CONDUCTED ON THE 11/11/1994




Figure 1. A 100% dot pattern used for the comparison of the dot pattern.




Figure 2. A 100% dot pattern used for the comparison of the dot pattern.




Figure 3. A 100% dot pattern used for the comparison of the dot pattern.




Figure 4. A 100% dot pattern used for the comparison of the dot pattern.




Figure 1. A 100% dot pattern used for the comparison of the dot pattern.




Figure 2. A 100% dot pattern used for the comparison of the dot pattern.




Figure 3. A 100% dot pattern used for the comparison of the dot pattern.






Figure 4. A 100% dot pattern used for the comparison of the dot pattern.




THE UNIVERSITY OF CHICAGO




THE UNIVERSITY OF CHICAGO




1. The first step is to identify the problem or question that needs to be answered. This involves understanding the context and the specific requirements of the task.




100







1000000




100







THESE ARE LAMINAR PROJECTIONS OF THE VOID SHOWN AT THE 15° ANGLE



THESE ARE LAMINAR PROJECTIONS OF THE VOID SHOWN AT THE 15° ANGLE











THESE ARE LAMINAR PROJECTIONS OF THE VOID SHOWN AT THE 15° ANGLE



THESE ARE LAMINAR PROJECTIONS OF THE VOID SHOWN AT THE 15° ANGLE

Figure 4

The particle positions through half of the void shown at various times in stereo projections looking down into the void at a 15° angle.






Fig. 1. Particle positions through the void at a 15° angle.




Fig. 2. Particle positions through the void at a 15° angle.




Fig. 3. Particle positions through the void at a 15° angle.












Fig. 4. Particle positions through the void at a 15° angle.

Figure 5

The particle positions through all of the void shown at various times in stereo projections looking down into the void at a 15° angle.






FIG. 1. a) Initial state; b) state after 100 time units.




FIG. 2. c) Initial state; d) state after 100 time units.

APPENDIX A

THE HYDRODYNAMIC EQUATIONS

The Particle-In-Cell method of Harlow¹ has been used by Mader^{2,3} to solve reactive two-dimensional fluid flow problems. We shall present a description of the equations used to solve the three-dimensional fluid flow problem. The equations are a straightforward extension of the EIC equations described by Mader².

The three-dimensional partial differential equations for nonviscous, nonconducting, compressible fluid flow are:

$$\begin{aligned} \rho \left[\frac{\partial \rho}{\partial t} + u \left(\frac{\partial \rho}{\partial x} \right) + v \left(\frac{\partial \rho}{\partial y} \right) + w \left(\frac{\partial \rho}{\partial z} \right) \right] &= -\rho \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} \right) \\ \rho \left[\frac{\partial u}{\partial t} + u \left(\frac{\partial u}{\partial x} \right) + v \left(\frac{\partial u}{\partial y} \right) + w \left(\frac{\partial u}{\partial z} \right) \right] &= -\frac{\partial p}{\partial x} \\ \rho \left[\frac{\partial v}{\partial t} + u \left(\frac{\partial v}{\partial x} \right) + v \left(\frac{\partial v}{\partial y} \right) + w \left(\frac{\partial v}{\partial z} \right) \right] &= -\frac{\partial p}{\partial y} \\ \rho \left[\frac{\partial w}{\partial t} + u \left(\frac{\partial w}{\partial x} \right) + v \left(\frac{\partial w}{\partial y} \right) + w \left(\frac{\partial w}{\partial z} \right) \right] &= -\frac{\partial p}{\partial z} \\ \rho \left[\frac{\partial E}{\partial t} + u \left(\frac{\partial E}{\partial x} \right) + v \left(\frac{\partial E}{\partial y} \right) + w \left(\frac{\partial E}{\partial z} \right) \right] &= -p \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} \right) \end{aligned}$$

In the PIC method, the dynamic quantities of the fluid are associated with a fixed (Eulerian) mesh of cells, and the fluid itself is represented by a set of mass points or particles resident in the cells. The conservation of mass is automatically satisfied by the particle model. Since the particles are not moved in Phase I, the transport terms in the momentum and energy equations may be dropped, resulting in the following set of equations.

$$\begin{aligned} \rho \frac{\partial u}{\partial t} &= - \frac{\partial p}{\partial x} \\ \rho \frac{\partial v}{\partial t} &= - \frac{\partial p}{\partial y} \\ \rho \frac{\partial w}{\partial t} &= - \frac{\partial p}{\partial z} \end{aligned} \quad \text{Momentum}$$


$$\rho \frac{\partial H}{\partial t} = - p \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} \right) \quad \text{Energy}$$

With artificial viscosity these equations become:


$$\begin{aligned} \rho \frac{\partial u}{\partial t} &= - \frac{\partial(p+q)}{\partial x} \\ \rho \frac{\partial v}{\partial t} &= - \frac{\partial(p+q)}{\partial y} \\ \rho \frac{\partial w}{\partial t} &= - \frac{\partial(p+q)}{\partial z} \\ - \rho \frac{\partial H}{\partial t} &= p \left[\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} \right] + \\ &+ \frac{\partial(uq)}{\partial x} - u \frac{\partial q}{\partial x} + \frac{\partial(vq)}{\partial y} - v \frac{\partial q}{\partial y} \\ &+ \frac{\partial(wq)}{\partial z} - w \frac{\partial q}{\partial z} \end{aligned}$$

Given some initial conditions, these equations are solved using finite-difference approximations.

Each cycle advancement is performed in three phases. In Phase I the cell quantities are updated using the above differential equations. In Phase II the particles are moved carrying with them the appropriate fractions of mass, momentum, and energy. Finally, in Phase III the amount of chemical reaction is calculated and the pressure and temperature of each cell are calculated using the HOF equation of state.



The problem is set up with a cubical lattice of 1728 cells (12 in each direction). Every cell has three indices (i,j,k) associated with it, each index having a range from zero to eleven. The (x, y, z) coordinates of the corner nearest the origin of cell ijk are then (iS, jS, kS) where S is the length of the side of a cell. The faces of each cell are numbered one through six.



The faces of cell ijk have indices

- face 1 i, j - 1/2, k
- face 2 i + 1/2, j, k
- face 3 i, j + 1/2, k
- face 4 i - 1/2, j, k

face 5 $i, j, k + 1/2$
face 6 $i, j, k - 1/2$

The particles are initially arranged in the same pattern in all cells. Their coordinates are such that no two particles in a cell have the same x, y , or z coordinates. For example, in cell $i = 0, j = 0, k = 0$, with unit side length, the sixteen particles have the following coordinates.

$(\frac{1}{17}, \frac{1}{17}, \frac{1}{17})$	$(\frac{5}{17}, \frac{3}{17}, \frac{27}{34})$	$(\frac{9}{17}, \frac{2}{17}, \frac{13}{38})$	$(\frac{13}{17}, \frac{1}{17}, \frac{35}{58})$
$(\frac{2}{17}, \frac{8}{17}, \frac{21}{34})$	$(\frac{6}{17}, \frac{7}{17}, \frac{6}{17})$	$(\frac{10}{17}, \frac{6}{17}, \frac{57}{58})$	$(\frac{14}{17}, \frac{5}{17}, \frac{5}{58})$
$(\frac{3}{17}, \frac{12}{17}, \frac{3}{17})$	$(\frac{7}{17}, \frac{11}{17}, \frac{21}{34})$	$(\frac{11}{17}, \frac{10}{17}, \frac{27}{38})$	$(\frac{15}{17}, \frac{2}{17}, \frac{53}{58})$
$(\frac{4}{17}, \frac{16}{17}, \frac{25}{34})$	$(\frac{8}{17}, \frac{15}{17}, \frac{8}{17})$	$(\frac{12}{17}, \frac{14}{17}, \frac{55}{58})$	$(\frac{16}{17}, \frac{13}{17}, \frac{13}{58})$

To begin the problem, the cell quantities (mass, velocities, pressure, temperature, internal energy, and mass fraction) are set to initial values and the particle mass is set to $1/16$ of the mass of its resident cell.

PHASE I

Phase I is not performed for empty cells nor for piston boundary cells.

A. Calculate the pressures on the six faces of each cell.

1. If a cell face is on a boundary, its pressure is set equal to the cell pressure.
2. If a neighbor cell is empty, the pressure on that face is set to zero.
3. Otherwise:

$$\begin{aligned} P_{i,j,k}^1 &= \frac{1}{2} (P_{i,j,k} + P_{i,j-1,k}) && \text{face 1 of cell } i,j,k \\ P_{i,j,k}^2 &= \frac{1}{2} (P_{i,j,k} + P_{i+1,j,k}) && \text{face 2 of cell } i,j,k \\ P_{i,j,k}^3 &= \frac{1}{2} (P_{i,j,k} + P_{i,j+1,k}) && \text{face 3 of cell } i,j,k \\ P_{i,j,k}^4 &= \frac{1}{2} (P_{i,j,k} + P_{i-1,j,k}) && \text{face 4 of cell } i,j,k \\ P_{i,j,k}^5 &= \frac{1}{2} (P_{i,j,k} + P_{i,j,k+1}) && \text{face 5 of cell } i,j,k \\ P_{i,j,k}^6 &= \frac{1}{2} (P_{i,j,k} + P_{i,j,k-1}) && \text{face 6 of cell } i,j,k \end{aligned}$$

B. Calculate the viscosities on the six faces of each cell.

1. If a neighbor cell is empty, the viscosity on that face is set to zero.
2. If the viscosity is negative, it is set to zero.
3. If a cell face is on a boundary, its viscosity is set to zero.
4. For the problems run on the $12 \times 12 \times 12$ cell lattice, $K = 1$ was sufficient. It may be necessary to go to higher or variable K 's in problems of greater resolution.

$$\begin{aligned}
Q_{1,j,k}^1 &= \frac{K M_{1,j,k} M_{1,j-1,k} (U_{y,i,j-1,k} - U_{y,i,j,k})}{2(M_{1,j,k} + M_{1,j-1,k}) S^3} \\
Q_{1,j,k}^2 &= \frac{K M_{1,j,k} M_{1,j+1,k} (U_{x,i,j,k} - U_{x,i+1,j,k})}{2(M_{1,j,k} + M_{1,j+1,k}) S^3} \\
Q_{1,j,k}^3 &= \frac{K V_{1,j,k} M_{1,j+1,k} (U_{y,i,j,k} - U_{y,i+1,k})}{2(M_{1,j,k} + M_{1,j+1,k}) S^3} \\
Q_{1,j,k}^4 &= \frac{K M_{1,j,k} M_{1,j-1,k} (U_{x,i-1,j,k} - U_{x,i,j,k})}{2(M_{1,j,k} + M_{1,j-1,k}) S^3} \\
Q_{1,j,k}^5 &= \frac{K M_{1,j,k} M_{1,j,k+1} (U_{z,i,j,k} - U_{z,i,j,k+1})}{2(M_{1,j,k} + M_{1,j,k+1}) S^3} \\
Q_{1,j,k}^6 &= \frac{K M_{1,j,k} M_{1,j,k-1} (U_{z,i,j,k-1} - U_{z,i,j,k})}{2(M_{1,j,k} + M_{1,j,k-1}) S^3}
\end{aligned}$$

C. Tentative velocities are calculated according to the momentum equations above.

$$\begin{aligned}
\bar{U}_{x,i,j,k} &= U_{x,i,j,k} + \frac{\Delta t S^2}{M_{1,j,k}} \left(P_{1,j,k}^4 - P_{1,j,k}^2 + Q_{1,j,k}^4 - Q_{1,j,k}^2 \right) \\
\bar{U}_{y,i,j,k} &= U_{y,i,j,k} + \frac{\Delta t S^2}{M_{1,j,k}} \left(P_{1,j,k}^1 - P_{1,j,k}^3 + Q_{1,j,k}^1 - Q_{1,j,k}^3 \right) \\
\bar{U}_{z,i,j,k} &= U_{z,i,j,k} + \frac{\Delta t S^2}{M_{1,j,k}} \left(P_{1,j,k}^6 - P_{1,j,k}^5 + Q_{1,j,k}^6 - Q_{1,j,k}^5 \right)
\end{aligned}$$

D. Calculate tentative cell internal energies.

1. Calculate velocity averages. If a neighbor cell is empty, the average velocity for that face is $\frac{1}{2} (\bar{U}_{1,j,k} + U_{1,j,k})$.

$$\begin{aligned}
U_{x,i,j,k}^4 &= \frac{1}{4} (\bar{U}_{x,i,j,k} + U_{x,i,j,k} + \bar{U}_{x,i-1,j,k} + U_{x,i-1,j,k}) \\
U_{x,i,j,k}^2 &= \frac{1}{4} (\bar{U}_{x,i,j,k} + U_{x,i,j,k} + \bar{U}_{x,i+1,j,k} + U_{x,i+1,j,k}) \\
U_{y,i,j,k}^1 &= \frac{1}{4} (\bar{U}_{y,i,j,k} + U_{y,i,j,k} + \bar{U}_{y,i,j-1,k} + U_{y,i,j-1,k}) \\
U_{y,i,j,k}^3 &= \frac{1}{4} (\bar{U}_{y,i,j,k} + U_{y,i,j,k} + \bar{U}_{y,i,j+1,k} + U_{y,i,j+1,k}) \\
U_{z,i,j,k}^6 &= \frac{1}{4} (\bar{U}_{z,i,j,k} + U_{z,i,j,k} + \bar{U}_{z,i,j,k-1} + U_{z,i,j,k-1}) \\
U_{z,i,j,k}^5 &= \frac{1}{4} (\bar{U}_{z,i,j,k} + U_{z,i,j,k} + \bar{U}_{z,i,j,k+1} + U_{z,i,j,k+1})
\end{aligned}$$

2. The tentative cell energy is then

$$\begin{aligned}
\bar{I}_{1,j,k} &= I_{1,j,k} + \frac{\Delta t S^2}{M_{1,j,k}} \left\{ P_{1,j,k}^4 (U_{x,i,j,k}^4 - U_{x,i,j,k}^2) \right. \\
&\quad + U_{y,i,j,k}^1 - U_{y,i,j,k}^3 + U_{z,i,j,k}^6 - U_{z,i,j,k}^5 \\
&\quad + Q_{1,j,k}^4 U_{x,i,j,k}^4 - Q_{1,j,k}^2 U_{x,i,j,k}^2 \\
&\quad + Q_{1,j,k}^1 U_{y,i,j,k}^1 - Q_{1,j,k}^3 U_{y,i,j,k}^3 + Q_{1,j,k}^6 U_{z,i,j,k}^6 \\
&\quad - Q_{1,j,k}^5 U_{z,i,j,k}^5 \\
&\quad - \frac{\bar{U}_{x,i,j,k} + U_{x,i,j,k}}{2} (Q_{1,j,k}^4 - Q_{1,j,k}^2) \\
&\quad - \frac{\bar{U}_{y,i,j,k} + U_{y,i,j,k}}{2} (Q_{1,j,k}^1 - Q_{1,j,k}^3) \\
&\quad \left. - \frac{\bar{U}_{z,i,j,k} + U_{z,i,j,k}}{2} (Q_{1,j,k}^6 - Q_{1,j,k}^5) \right\}
\end{aligned}$$

E. Calculate total cell energy and momenta.

$$\bar{E}_{i,j,k} = E_{i,j,k} \left[\bar{I}_{i,j,k} + \frac{1}{2} (\bar{U}_{x,i,j,k}^2 + \bar{U}_{y,i,j,k}^2 + \bar{U}_{z,i,j,k}^2) \right]$$

$$\bar{X}_{x,i,j,k} = N_{i,j,k} \bar{U}_{x,i,j,k}$$

$$\bar{X}_{y,i,j,k} = N_{i,j,k} \bar{U}_{y,i,j,k}$$

$$\bar{X}_{z,i,j,k} = N_{i,j,k} \bar{U}_{z,i,j,k}$$

PHASE II

For the movement of the particles, the velocity is determined by a method of weighting. A cube, the size of a cell, is imagined with the particle at its center. In general, this cube intersects eight cells with a certain fraction of its volume in each. The tentative velocities of the overlapped cells, weighted by their respective fractions of the volume, are added together to form the particle velocity. Those parts of the cube that fall outside the boundaries or intersect an empty cell are not used in the weighting.

The position of the particles in the cell determines which neighboring cells are used in the weighting. For a particle with coordinates (x, y, z) residing in cell ijk (i.e., $i \leq x/S \leq i+1$; $j \leq y/S \leq j+1$; and $k \leq z/S \leq k+1$) the eight different cases are:

1. $i \leq x/S \leq i+1/2$ 3. $i \leq x/S \leq i+1/2$ 5. $i+1/2 \leq x/S \leq i+1$
 $j \leq y/S \leq j+1/2$ $j+1/2 \leq y/S \leq j+1$ $j \leq y/S \leq j+1/2$
 $k \leq z/S \leq k+1/2$ $k+1/2 \leq z/S \leq k+1$ $k \leq z/S \leq k+1/2$
2. $i \leq x/S \leq i+1/2$ 4. $i \leq x/S \leq i+1/2$ 6. $i+1/2 \leq x/S \leq i+1$
 $j+1/2 \leq y/S \leq j+1$ $j \leq y/S \leq j+1/2$ $j+1/2 \leq y/S \leq j+1$
 $k \leq z/S \leq k+1/2$ $k+1/2 \leq z/S \leq k+1$ $k \leq z/S \leq k+1/2$

7. $i+1/2 \leq x/S \leq i+1$ 8. $i+1/2 \leq x/S \leq i+1$
 $j+1/2 \leq y/S \leq j+1$ $j \leq y/S \leq j+1/2$
 $k+1/2 \leq z/S \leq k+1$ $k+1/2 \leq z/S \leq k+1$

The weighted particle velocities for each of these cases are as follows $[(x, y, z)]$ the coordinates of the particle in cell ijk with no faces on a boundary].

1. $V_1 = (1S + S/2 - x) (jS + S/2 - y) (kS + S/2 - z)$
 $V_2 = (1S + S/2 - x) (y + S/2 - jS) (kS + S/2 - z)$
 $V_3 = (1S + S/2 - x) (y + S/2 - jS) (z + S/2 - kS)$
 $V_4 = (1S + S/2 - x) (jS + S/2 - y) (z + S/2 - kS)$
 $V_5 = (x + S/2 - 1S) (jS + S/2 - y) (kS + S/2 - z)$
 $V_6 = (x + S/2 - 1S) (y + S/2 - jS) (kS + S/2 - z)$
 $V_7 = (x + S/2 - 1S) (y + S/2 - jS) (z + S/2 - kS)$
 $V_8 = (x + S/2 - 1S) (jS + S/2 - y) (z + S/2 - kS)$

$$U_x^1 = \left(V_1 \bar{U}_{x,i-1,j-1,k-1} + V_2 \bar{U}_{x,i-1,j,k-1} + V_3 \bar{U}_{x,i-1,j,k} \right. \\
+ V_4 \bar{U}_{x,i-1,j-1,k} + V_5 \bar{U}_{x,i,j-1,k-1} + V_6 \bar{U}_{x,i,j,k-1} \\
+ V_7 \bar{U}_{x,i,j,k} + V_8 \bar{U}_{x,i,j-1,k} \Big) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 + V_7 + V_8)$$

$$U_y^1 = \left(V_1 \bar{U}_{y,i-1,j-1,k-1} + V_2 \bar{U}_{y,i-1,j,k-1} + V_3 \bar{U}_{y,i-1,j,k} \right. \\
+ V_4 \bar{U}_{y,i-1,j-1,k} + V_5 \bar{U}_{y,i,j-1,k-1} + V_6 \bar{U}_{y,i,j,k-1} \\
+ V_7 \bar{U}_{y,i,j,k} + V_8 \bar{U}_{y,i,j-1,k} \Big) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 + V_7 + V_8)$$

$$u_z^1 = \left(V_1 \bar{u}_{z,i-1,j-1,k-1} + V_2 \bar{u}_{z,i-1,j,k-1} + V_3 \bar{u}_{z,i-1,j,k} \right. \\ \left. + V_4 \bar{u}_{z,i-1,j-1,k} + V_5 \bar{u}_{z,i,j-1,k-1} + V_6 \bar{u}_{z,i,j,k-1} \right. \\ \left. + V_7 \bar{u}_{z,i,j,k} + V_8 \bar{u}_{z,i,j-1,k} \right) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 \\ + V_7 + V_8)$$

$$2. \quad V_1 = (1S + S/2 - x) (jS + 3S/2 - y) (kS + S/2 - z) \\ V_2 = (1S + S/2 - x) (y - S/2 - jS) (kS + S/2 - z) \\ V_3 = (1S + S/2 - x) (y - S/2 - jS) (z + S/2 - kS) \\ V_4 = (1S + S/2 - x) (jS + 3S/2 - y) (z + S/2 - kS) \\ V_5 = (x + S/2 - iS) (jS + 3S/2 - y) (kS + S/2 - z) \\ V_6 = (x + S/2 - iS) (y - S/2 - jS) (kS + S/2 - z) \\ V_7 = (x + S/2 - iS) (y - S/2 - jS) (z + S/2 - kS) \\ V_8 = (x + S/2 - iS) (jS + 3S/2 - y) (z + S/2 - kS)$$

$$u_x^1 = \left(V_1 \bar{u}_{x,i-1,j,k-1} + V_2 \bar{u}_{x,i-1,j+1,k-1} + V_3 \bar{u}_{x,i-1,j+1,k} \right. \\ \left. + V_4 \bar{u}_{x,i-1,j,k} + V_5 \bar{u}_{x,i,j,k-1} + V_6 \bar{u}_{x,i,j+1,k-1} \right. \\ \left. + V_7 \bar{u}_{x,i,j+1,k} + V_8 \bar{u}_{x,i,j,k} \right) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 \\ + V_7 + V_8)$$

$$u_y^1 = \left(V_1 \bar{u}_{y,i-1,j,k-1} + V_2 \bar{u}_{y,i-1,j+1,k-1} + V_3 \bar{u}_{y,i-1,j+1,k} \right. \\ \left. + V_4 \bar{u}_{y,i-1,j,k} + V_5 \bar{u}_{y,i,j,k-1} + V_6 \bar{u}_{y,i,j+1,k-1} \right. \\ \left. + V_7 \bar{u}_{y,i,j+1,k} + V_8 \bar{u}_{y,i,j,k} \right) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 \\ + V_7 + V_8)$$

$$u_z^1 = \left(V_1 \bar{u}_{z,i-1,j,k-1} + V_2 \bar{u}_{z,i-1,j+1,k-1} + V_3 \bar{u}_{z,i-1,j+1,k} \right. \\ \left. + V_4 \bar{u}_{z,i-1,j,k-1} + V_5 \bar{u}_{z,i,j,k-1} + V_6 \bar{u}_{z,i,j+1,k-1} \right. \\ \left. + V_7 \bar{u}_{z,i,j+1,k} + V_8 \bar{u}_{z,i,j,k} \right) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 \\ + V_7 + V_8)$$

$$3. \quad V_1 = (1S + S/2 - x) (jS + 3S/2 - y) (kS + 3S/2 - z) \\ V_2 = (1S + S/2 - x) (y - S/2 - jS) (kS + 3S/2 - z) \\ V_3 = (1S + S/2 - x) (y - S/2 - jS) (z - S/2 - kS) \\ V_4 = (1S + S/2 - x) (jS + 3S/2 - y) (z - S/2 - kS) \\ V_5 = (x + S/2 - iS) (jS + 3S/2 - y) (kS + 3S/2 - z) \\ V_6 = (x + S/2 - iS) (y - S/2 - jS) (kS + 3S/2 - z) \\ V_7 = (x + S/2 - iS) (y - S/2 - jS) (z - S/2 - kS) \\ V_8 = (x + S/2 - iS) (jS + 3S/2 - y) (z - S/2 - kS)$$

$$u_x^1 = \left(V_1 \bar{u}_{x,i-1,j,k} + V_2 \bar{u}_{x,i-1,j+1,k} + V_3 \bar{u}_{x,i-1,j+1,k+1} \right. \\ \left. + V_4 \bar{u}_{x,i-1,j,k+1} + V_5 \bar{u}_{x,i,j,k} + V_6 \bar{u}_{x,i,j+1,k} \right. \\ \left. + V_7 \bar{u}_{x,i,j+1,k+1} + V_8 \bar{u}_{x,i,j,k+1} \right) / (V_1 + V_2 + V_3 + V_4 + V_5 \\ + V_6 + V_7 + V_8)$$

$$u_y^1 = \left(V_1 \bar{u}_{y,i-1,j,k} + V_2 \bar{u}_{y,i-1,j+1,k} + V_3 \bar{u}_{y,i-1,j+1,k+1} \right. \\ \left. + V_4 \bar{u}_{y,i-1,j,k+1} + V_5 \bar{u}_{y,i,j,k} + V_6 \bar{u}_{y,i,j+1,k} \right. \\ \left. + V_7 \bar{u}_{y,i,j+1,k+1} + V_8 \bar{u}_{y,i,j,k+1} \right) / (V_1 + V_2 + V_3 + V_4 + V_5 \\ + V_6 + V_7 + V_8)$$

$$U_z^1 = (V_1 \bar{U}_{z,i-1,j,k} + V_2 \bar{U}_{z,i-1,j+1,k} + V_3 \bar{U}_{z,i-1,j,k+1} + V_4 \bar{U}_{z,i-1,j,k+1} + V_5 \bar{U}_{z,i,j-1,k} + V_6 \bar{U}_{z,i,j,k} + V_7 \bar{U}_{z,i,j,k+1} + V_8 \bar{U}_{z,i,j-1,k+1}) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 + V_7 + V_8)$$

$$\begin{aligned} V_1 &= (1S + 3S/2 - x) (jS + S/2 - y) (kS + S/2 - z) \\ V_2 &= (1S + 3S/2 - x) (y + S/2 - jS) (kS + S/2 - z) \\ V_3 &= (1S + 3S/2 - x) (y + S/2 - jS) (z + S/2 - kS) \\ V_4 &= (1S + 3S/2 - x) (jS + S/2 - y) (z + S/2 - kS) \\ V_5 &= (x - S/2 - iS) (jS + S/2 - y) (kS + S/2 - z) \\ V_6 &= (x - S/2 - iS) (y + S/2 - jS) (kS + S/2 - z) \\ V_7 &= (x - S/2 - iS) (y + S/2 - jS) (z + S/2 - kS) \\ V_8 &= (x - S/2 - iS) (jS + S/2 - y) (z + S/2 - kS) \end{aligned}$$

$$U_x^1 = (V_1 \bar{U}_{x,i,j-1,k-1} + V_2 \bar{U}_{x,i,j,k-1} + V_3 \bar{U}_{x,i,j,k} + V_4 \bar{U}_{x,i,j-1,k} + V_5 \bar{U}_{x,i+1,j-1,k-1} + V_6 \bar{U}_{x,i+1,j,k-1} + V_7 \bar{U}_{x,i+1,j,k} + V_8 \bar{U}_{x,i+1,j-1,k}) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 + V_7 + V_8)$$

$$U_y^1 = (V_1 \bar{U}_{y,i,j-1,k-1} + V_2 \bar{U}_{y,i,j,k-1} + V_3 \bar{U}_{y,i,j,k} + V_4 \bar{U}_{y,i,j-1,k} + V_5 \bar{U}_{y,i+1,j-1,k-1} + V_6 \bar{U}_{y,i+1,j,k-1} + V_7 \bar{U}_{y,i+1,j,k} + V_8 \bar{U}_{y,i+1,j-1,k}) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 + V_7 + V_8)$$

-49-

$$U_z^1 = (V_1 \bar{U}_{z,i-1,j,k} + V_2 \bar{U}_{z,i-1,j+1,k} + V_3 \bar{U}_{z,i-1,j,k+1} + V_4 \bar{U}_{z,i-1,j,k+1} + V_5 \bar{U}_{z,i,j-1,k} + V_6 \bar{U}_{z,i,j,k} + V_7 \bar{U}_{z,i,j,k+1} + V_8 \bar{U}_{z,i,j-1,k+1}) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 + V_7 + V_8)$$

$$\begin{aligned} V_1 &= (1S + S/2 - x) (jS + S/2 - y) (kS + 3S/2 - z) \\ V_2 &= (1S + S/2 - x) (y + S/2 - jS) (kS + 3S/2 - z) \\ V_3 &= (1S + S/2 - x) (y + S/2 - jS) (z - S/2 - kS) \\ V_4 &= (1S + S/2 - x) (jS + S/2 - y) (z - S/2 - kS) \\ V_5 &= (x + S/2 - iS) (jS + S/2 - y) (kS + 3S/2 - z) \\ V_6 &= (x + S/2 - iS) (y + S/2 - jS) (kS + 3S/2 - z) \\ V_7 &= (x + S/2 - iS) (y + S/2 - jS) (z - S/2 - kS) \\ V_8 &= (x + S/2 - iS) (jS + S/2 - y) (z - S/2 - kS) \end{aligned}$$

$$U_x^1 = (V_1 \bar{U}_{x,i,j-1,k} + V_2 \bar{U}_{x,i,j,k} + V_3 \bar{U}_{x,i,j,k+1} + V_4 \bar{U}_{x,i,j-1,k+1} + V_5 \bar{U}_{x,i,j-1,k} + V_6 \bar{U}_{x,i,j,k} + V_7 \bar{U}_{x,i,j,k+1} + V_8 \bar{U}_{x,i,j-1,k+1}) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 + V_7 + V_8)$$

$$U_y^1 = (V_1 \bar{U}_{y,i,j-1,k} + V_2 \bar{U}_{y,i,j,k} + V_3 \bar{U}_{y,i,j,k+1} + V_4 \bar{U}_{y,i,j-1,k+1} + V_5 \bar{U}_{y,i,j-1,k} + V_6 \bar{U}_{y,i,j,k} + V_7 \bar{U}_{y,i,j,k+1} + V_8 \bar{U}_{y,i,j-1,k+1}) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 + V_7 + V_8)$$

-48-

$$u_z^1 = \left(v_1 \bar{u}_{z,i,j,k-1} + v_2 \bar{u}_{z,i,j+1,k-1} + v_3 \bar{u}_{z,i,j+1,k} \right. \\ \left. + v_4 \bar{u}_{z,i,j,k} + v_5 \bar{u}_{z,i+1,j,k-1} + v_6 \bar{u}_{z,i+1,j+1,k-1} \right. \\ \left. + v_7 \bar{u}_{z,i+1,j+1,k} + v_8 \bar{u}_{z,i+1,j,k} \right) / (v_1 + v_2 + v_3 + v_4 + v_5 \\ + v_6 + v_7 + v_8)$$

$$7. \quad \begin{aligned} v_1 &= (1S + 3S/2 - x) (jS + 3S/2 - y) (kS + 3S/2 - z) \\ v_2 &= (1S + 3S/2 - x) (y - S/2 - jS) (kS + 3S/2 - z) \\ v_3 &= (1S + 3S/2 - x) (y - S/2 - jS) (z - S/2 - kS) \\ v_4 &= (1S + 3S/2 - x) (jS + 3S/2 - y) (z - S/2 - kS) \\ v_5 &= (x - S/2 - 1S) (jS + 3S/2 - y) (kS + 3S/2 - z) \\ v_6 &= (x - S/2 - 1S) (y - S/2 - jS) (kS + 3S/2 - z) \\ v_7 &= (x - S/2 - 1S) (y - S/2 - jS) (z - S/2 - kS) \\ v_8 &= (x - S/2 - 1S) (jS + 3S/2 - y) (z - S/2 - kS) \end{aligned}$$

$$u_x^1 = \left(v_1 \bar{u}_{x,i,j,k} + v_2 \bar{u}_{x,i,j+1,k} + v_3 \bar{u}_{x,i,j+1,k+1} \right. \\ \left. + v_4 \bar{u}_{x,i,j,k+1} + v_5 \bar{u}_{x,i+1,j,k} + v_6 \bar{u}_{x,i+1,j+1,k} \right. \\ \left. + v_7 \bar{u}_{x,i+1,j+1,k+1} + v_8 \bar{u}_{x,i+1,j,k+1} \right) / (v_1 + v_2 + v_3 + v_4 \\ + v_5 + v_6 + v_7 + v_8)$$

$$u_y^1 = \left(v_1 \bar{u}_{y,i,j,k} + v_2 \bar{u}_{y,i,j+1,k} + v_3 \bar{u}_{y,i,j+1,k+1} \right. \\ \left. + v_4 \bar{u}_{y,i,j,k+1} + v_5 \bar{u}_{y,i+1,j,k} + v_6 \bar{u}_{y,i+1,j+1,k} \right. \\ \left. + v_7 \bar{u}_{y,i+1,j+1,k+1} + v_8 \bar{u}_{y,i+1,j,k+1} \right) / (v_1 + v_2 + v_3 + v_4 \\ + v_5 + v_6 + v_7 + v_8)$$

-51-

$$u_z^1 = \left(v_1 \bar{u}_{z,i,j-1,k-1} + v_2 \bar{u}_{z,i,j,k-1} + v_3 \bar{u}_{z,i,j,k} \right. \\ \left. + v_4 \bar{u}_{z,i,j-1,k} + v_5 \bar{u}_{z,i+1,j-1,k-1} + v_6 \bar{u}_{z,i+1,j,k-1} \right. \\ \left. + v_7 \bar{u}_{z,i+1,j,k} + v_8 \bar{u}_{z,i+1,j-1,k} \right) / (v_1 + v_2 + v_3 + v_4 + v_5 \\ + v_6 + v_7 + v_8)$$

$$6. \quad \begin{aligned} v_1 &= (1S + 3S/2 - x) (jS + 3S/2 - y) (kS + S/2 - z) \\ v_2 &= (1S + 3S/2 - x) (y - S/2 - jS) (kS + S/2 - z) \\ v_3 &= (1S + 3S/2 - x) (y - S/2 - jS) (z + S/2 - kS) \\ v_4 &= (1S + 3S/2 - x) (jS + 3S/2 - y) (z + S/2 - kS) \\ v_5 &= (x - S/2 - 1S) (jS + 3S/2 - y) (kS + S/2 - z) \\ v_6 &= (x - S/2 - 1S) (y - S/2 - jS) (kS + S/2 - z) \\ v_7 &= (x - S/2 - 1S) (y - S/2 - jS) (z + S/2 - kS) \\ v_8 &= (x - S/2 - 1S) (jS + 3S/2 - y) (z + S/2 - kS) \end{aligned}$$

$$u_x^1 = \left(v_1 \bar{u}_{x,i,j,k-1} + v_2 \bar{u}_{x,i,j+1,k-1} + v_3 \bar{u}_{x,i,j+1,k} \right. \\ \left. + v_4 \bar{u}_{x,i,j,k} + v_5 \bar{u}_{x,i+1,j,k-1} + v_6 \bar{u}_{x,i+1,j+1,k-1} \right. \\ \left. + v_7 \bar{u}_{x,i+1,j+1,k} + v_8 \bar{u}_{x,i+1,j,k} \right) / (v_1 + v_2 + v_3 + v_4 + v_5 \\ + v_6 + v_7 + v_8)$$

$$u_y^1 = \left(v_1 \bar{u}_{y,i,j,k-1} + v_2 \bar{u}_{y,i,j+1,k-1} + v_3 \bar{u}_{y,i,j+1,k} \right. \\ \left. + v_4 \bar{u}_{y,i,j,k} + v_5 \bar{u}_{y,i+1,j,k-1} + v_6 \bar{u}_{y,i+1,j+1,k-1} \right. \\ \left. + v_7 \bar{u}_{y,i+1,j+1,k} + v_8 \bar{u}_{y,i+1,j,k} \right) / (v_1 + v_2 + v_3 + v_4 + v_5 \\ + v_6 + v_7 + v_8)$$

-50-

$$U_z^1 = \left(V_1 \bar{U}_{x,i,j,k} + V_2 \bar{U}_{x,i,j+1,k} + V_3 \bar{U}_{x,i,j+1,k+1} + V_4 \bar{U}_{x,i,j,k+1} + V_5 \bar{U}_{x,i+1,j,k} + V_6 \bar{U}_{x,i+1,j+1,k} + V_7 \bar{U}_{x,i+1,j+1,k+1} + V_8 \bar{U}_{x,i+1,j,k+1} \right) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 + V_7 + V_8)$$

$$\begin{aligned} 8. \quad V_1 &= (1S + 3S/2 - x) (jS + S/2 - y) (kS + 3S/2 - z) \\ V_2 &= (1S + 3S/2 - x) (y + S/2 - jS) (kS + 3S/2 - z) \\ V_3 &= (1S + 3S/2 - x) (y + S/2 - jS) (z - S/2 - kS) \\ V_4 &= (1S + 3S/2 - x) (jS + S/2 - y) (z - S/2 - kS) \\ V_5 &= (x - S/2 - jS) (jS + S/2 - y) (kS + 3S/2 - z) \\ V_6 &= (x - S/2 - jS) (y + S/2 - jS) (kS + 3S/2 - z) \\ V_7 &= (x - S/2 - jS) (y + S/2 - jS) (z - S/2 - kS) \\ V_8 &= (x - S/2 - jS) (jS + S/2 - y) (z - S/2 - kS) \end{aligned}$$

$$U_x^1 = \left(V_1 \bar{U}_{x,i,j-1,k} + V_2 \bar{U}_{x,i,j,k} + V_3 \bar{U}_{x,i,j,k+1} + V_4 \bar{U}_{x,i,j-1,k+1} + V_5 \bar{U}_{x,i+1,j-1,k} + V_6 \bar{U}_{x,i+1,j,k} + V_7 \bar{U}_{x,i+1,j,k+1} + V_8 \bar{U}_{x,i+1,j-1,k+1} \right) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 + V_7 + V_8)$$

$$U_y^1 = \left(V_1 \bar{U}_{y,i,j-1,k} + V_2 \bar{U}_{y,i,j,k} + V_3 \bar{U}_{y,i,j,k+1} + V_4 \bar{U}_{y,i,j-1,k+1} + V_5 \bar{U}_{y,i+1,j-1,k} + V_6 \bar{U}_{y,i+1,j,k} + V_7 \bar{U}_{y,i+1,j,k+1} + V_8 \bar{U}_{y,i+1,j-1,k+1} \right) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 + V_7 + V_8)$$

$$U_z^1 = \left(V_1 \bar{U}_{z,i,j-1,k} + V_2 \bar{U}_{z,i,j,k} + V_3 \bar{U}_{z,i,j,k+1} + V_4 \bar{U}_{z,i,j-1,k+1} + V_5 \bar{U}_{z,i+1,j-1,k} + V_6 \bar{U}_{z,i+1,j,k} + V_7 \bar{U}_{z,i+1,j,k+1} + V_8 \bar{U}_{z,i+1,j-1,k+1} \right) / (V_1 + V_2 + V_3 + V_4 + V_5 + V_6 + V_7 + V_8)$$

The new position of the particle is calculated from these velocities.

$$x' = x + U_x^1 \Delta t$$

$$y' = y + U_y^1 \Delta t$$

$$z' = z + U_z^1 \Delta t$$

If a particle moves across a boundary out of the range of calculation, it is removed from the calculation by setting its mass to zero. The appropriate fractions of energy and momenta are subtracted from the cell it left.

If a particle (of mass m) moves from one cell (ijk) to another (lmn), the cell quantities are adjusted as follows.

$$\begin{aligned} H_{i,j,k}^1 &= H_{i,j,k} - m \\ H_{l,m,n}^1 &= H_{l,m,n} + m \end{aligned}$$

$$\begin{aligned} \bar{X}_{i,j,k}^1 &= \bar{X}_{i,j,k} - m \bar{U}_{x,i,j,k} \\ \bar{Y}_{i,j,k}^1 &= \bar{Y}_{i,j,k} - m \bar{U}_{y,i,j,k} \\ \bar{Z}_{i,j,k}^1 &= \bar{Z}_{i,j,k} - m \bar{U}_{z,i,j,k} \end{aligned}$$

$$\begin{aligned} \bar{X}_{l,m,n}^1 &= \bar{X}_{l,m,n} + m \bar{U}_{x,i,j,k} \\ \bar{Y}_{l,m,n}^1 &= \bar{Y}_{l,m,n} + m \bar{U}_{y,i,j,k} \\ \bar{Z}_{l,m,n}^1 &= \bar{Z}_{l,m,n} + m \bar{U}_{z,i,j,k} \end{aligned}$$

$$\begin{aligned}
 E_{i,j,k}^i &= E_{i,j,k} - \frac{m}{K_{i,j,k}} E_{i,j,k} \\
 E_{i,m,n}^i &= E_{i,m,n} + \frac{m}{K_{i,j,k}} E_{i,j,k} \\
 W_{i,m,n}^i &= \frac{H_{i,m,n} W_{i,m,n} + m W_{i,j,k}}{H_{i,m,n} + m}
 \end{aligned}$$

PHASE III

A. The velocities and internal energy are calculated from the momenta and total energy.

$$\begin{aligned}
 U_{x,i,j,k} &= \frac{\bar{X}_{x,i,j,k}}{M_{i,j,k}} \\
 U_{y,i,j,k} &= \frac{\bar{X}_{y,i,j,k}}{M_{i,j,k}} \\
 U_{z,i,j,k} &= \frac{\bar{X}_{z,i,j,k}}{M_{i,j,k}} \\
 I_{i,j,k} &= \frac{\bar{E}_{i,j,k}}{M_{i,j,k}} - \frac{1}{2} (U_{x,i,j,k}^2 + U_{y,i,j,k}^2 + U_{z,i,j,k}^2)
 \end{aligned}$$

B. Chemical reaction is performed according to the Arrhenius rate law.²

$$W_{i,j,k}^i = W_{i,j,k} \left[1 - (z) (\Delta t) e^{-E_a/R T_{i,j,k}} \right]$$

C. The equation of state used is the NKN equation of state.^{2,3} Given the specific volume, internal energy, and mass fraction of a cell, NKN calculates the temperature and pressure. For specific volumes less than the initial value, the pressure and temperature are set to their initial

values. If the mass of a cell is zero, the pressure and temperature are also set to zero.

D. The total time is increased by Δt , and the calculation begins with Phase I.

THE NOMENCLATURE

Quantities pertaining to cell i,j,k

$E_{i,j,k}$	total energy
$I_{i,j,k}$	internal energy
$M_{i,j,k}$	mass
$P_{i,j,k}$	pressure
$p_{i,j,k}^f$	pressure on face f
$\zeta_{i,j,k}^f$	viscosity pressure on face f
l	length of cell side (2×10^{-3} cm)
$T_{i,j,k}$	temperature
$U_{x,i,j,k}$	x velocity
$U_{y,i,j,k}$	y velocity
$U_{z,i,j,k}$	z velocity
$W_{i,j,k}$	mass fraction of undecomposed explosive
$X_{x,i,j,k}$	momentum in x direction
$X_{y,i,j,k}$	momentum in y direction
$X_{z,i,j,k}$	momentum in z direction

NOMENCLATURE (continued)

Quantities pertaining to each particle

m	mass
u_x^1	x velocity
u_y^1	y velocity
u_z^1	z velocity
x	x coordinate
y	y coordinate
z	z coordinate
Δt	time increment (5×10^{-4} sec)
ρ	density

APPENDIX B

STEREO PLOTTING

We shall describe a technique for stereo plotting a set of points in three dimensions. This method was developed by K. Crandall and R. Lewis of the Los Alamos Scientific Laboratory.

From the direction of sight, three unit vectors are determined.

$$\begin{aligned} \vec{e}_1 & \text{ in the direction of the viewer's line of sight} \\ \vec{e}_2 & \text{ points from the right eye to the left eye} \\ \vec{e}_3 & = \vec{e}_1 \times \vec{e}_2 \end{aligned}$$

A plane is then constructed perpendicular to \vec{e}_1 through the point with the minimum scalar product with \vec{e}_1 ; i.e., if the plane is moved closer to the viewer, all the points are beyond it, and if it is moved away, there are points between it and the viewer. All the points are projected onto this plane to determine the size square which will contain all the points in the three-dimensional views. From this window size and the characteristics of the stereo viewer, one determines the distance, d , that the eyes should be from this plane. The position of the eyes is then found by joining a distance d in the $-\vec{e}_1$ direction from the center of the window and then a short distance along \vec{e}_2 and $-\vec{e}_2$ to separate the two eyes. The window size is then divided by d to get the window size at a unit distance from the eyes. A plane is constructed at this point perpendicular to \vec{e}_1 with a coordinate system (x', y') ; the x' axis in the $-\vec{e}_2$ direction and the y' axis in the \vec{e}_3 direction.

The stereo pictures are produced by taking one eye at a time, connecting it by a straight line to each point, finding the intersection with the plane a unit distance from the eyes, and determining the (x', y') coordinates of the point of intersection. These coordinates are then used to plot the point on the SC-4020.

The mathematical techniques are as follows:

Given $\vec{e}_1 = (e_{1x}, e_{1y}, e_{1z})$ and a set of points $\{\vec{r}_k\}$ in three dimensions, define

$$\vec{e}_2 = \frac{1}{\sqrt{e_{1x}^2 + e_{1y}^2}} (e_{1y}, -e_{1x}, 0) \quad (a)$$

$$\vec{e}_3 = \vec{e}_1 \times \vec{e}_2$$

If e_{3z} is negative, replace \vec{e}_2 by $-\vec{e}_2$ and \vec{e}_3 by $-\vec{e}_3$. This assures that \vec{e}_2 points from the right eye to the left and \vec{e}_3 points upward.

The position of the plane of projection is determined by finding the point \vec{r}_m which minimizes $\vec{r}_k \cdot \vec{e}_1$ (b). One then projects each of the points \vec{r}_k into this plane and determines their (x_k', y_k') coordinates.


$$x_k' = -\vec{e}_2 \cdot \left[(\vec{r}_m - \vec{r}_k) \cdot \vec{e}_1 \right] \vec{e}_1 + \vec{r}_k \quad (c)$$

$$y_k' = \vec{e}_3 \cdot \left[(\vec{r}_m - \vec{r}_k) \cdot \vec{e}_1 \right] \vec{e}_1 + \vec{r}_k \quad (d)$$

$$\text{Let } \bar{x} = \max_k \{x_k'\}, \quad \bar{y} = \max_k \{y_k'\}$$

$$\hat{x} = \min_k \{x_k'\} \text{ and } \hat{y} = \min_k \{y_k'\}$$

For a given stereo viewer there is an angular limit, α , to the field of vision. The distance, d , of the eyes from the plane of projection is determined by T , the width of the window, and α . The half-



window width at a unit distance from the eyes $v = \frac{T}{d}$ is a constant for each viewer. The eyes are separated by a distance $\frac{2st}{1+s}$ where the value of s is again dependent upon the stereo viewer. The distance, d , the eyes must be from the plane in order to see everything in the x' direction is $d \geq \frac{(\bar{x} - \hat{x})(1+s)}{v}$. To see everything in the y direction, $d \geq \frac{(\bar{y} - \hat{y})}{v}$. The larger of these two distances is taken as the distance of the eyes from the center of the window \vec{e}_1 . The size, T , of a side of the square window is then either $(1+s)(\bar{x} - \hat{x})$ or $(\bar{y} - \hat{y})$.

The vector to the center of the projection is

$$\vec{c} = (\vec{r}_m \cdot \vec{e}_1) \vec{e}_1 - \frac{(\bar{x} + \hat{x})}{2} \vec{e}_2 + \frac{(\bar{y} + \hat{y})}{2} \vec{e}_3, \quad (e)$$

the vector to the right eye is

$$\vec{r}_r = \vec{c} - d\vec{e}_1 - \frac{T}{1+s} \vec{e}_2, \quad (f)$$

and the one to the left is

$$\vec{r}_l = \vec{c} - d\vec{e}_1 + \frac{T}{1+s} \vec{e}_2. \quad (g)$$

The two pictures are then made on the 35-4020, plotting the view for the left eye first. The coordinates of the point of intersection of a line from an eye to a point \vec{r}_k and a plane a unit distance away from the eyes are

$$x_k = - \left(\frac{\vec{r}_k - \vec{r}}{(\vec{r}_k - \vec{r}) \cdot \vec{e}_1} - \vec{e}_1 \right) \cdot \vec{e}_2, \quad (h)$$

and

$$y_k = \left(\frac{\vec{r}_k - \vec{r}}{(\vec{r}_k - \vec{r}) \cdot \vec{e}_1} - \vec{e}_1 \right) \cdot \vec{e}_3, \quad (i)$$

where \vec{r} is a vector to one of the eyes. These coordinates (x_k, y_k) are then plotted using the MATH routine on SUMTECH with grid boundaries of $-x$ to x in both x and y directions.

For Figures 5 and 6 we used the following parameters.

$$v = 0.3$$

$$s = 0.25$$

$$\vec{c}_1 = (0, 0.9593, -0.25882) = (0, \cos 15^\circ, -\sin 15^\circ)$$

$$\vec{c}_2 = (-1, 0, 0)$$

$$\vec{c}_3 = (0, 0.25082, 0.9593)$$

The coding formulas for the equations above designated by letters are as follows.

(a) The components of \vec{e}_3 are

$$e_{3x} = e_{1y} e_{2z} - e_{1z} e_{2y}$$

$$e_{3y} = e_{1z} e_{2x} - e_{1x} e_{2z}$$

$$e_{3z} = e_{1x} e_{2y} - e_{1y} e_{2x}$$

$$(b) \quad r_{mx} e_{1x} + r_{my} e_{1y} + r_{mz} e_{1z} = r_{mx} e_{1x} + r_{ky} e_{1y} + r_{kz} e_{1z} \text{ for all } k.$$

$$(c) \text{ and } (d) \text{ Since } \vec{e}_2 \cdot \vec{e}_1 = \vec{e}_3 \cdot \vec{e}_1 = 0,$$

$$x'_k = -e_{2x} r_{kx} - e_{2y} r_{ky} - e_{2z} r_{kz}$$

$$y'_k = e_{3x} r_{kx} + e_{3y} r_{ky} + e_{3z} r_{kz}$$

$$(e) \quad c_x = (r_{mx} e_{1x} + r_{my} e_{1y} + r_{mz} e_{1z}) e_{1x} - \left(\frac{\bar{x} + \bar{z}}{2} \right) e_{2x} + \left(\frac{\bar{y} + \bar{y}}{2} \right) e_{3x}$$

$$c_y = (r_{mx} e_{1x} + r_{my} e_{1y} + r_{mz} e_{1z}) e_{1y} - \left(\frac{\bar{x} + \bar{z}}{2} \right) e_{2y} + \left(\frac{\bar{y} + \bar{y}}{2} \right) e_{3y}$$

$$c_z = (r_{mx} e_{1x} + r_{my} e_{1y} + r_{mz} e_{1z}) e_{1z} - \left(\frac{\bar{x} + \bar{z}}{2} \right) e_{2z} + \left(\frac{\bar{y} + \bar{y}}{2} \right) e_{3z}$$

$$(f) \quad Y_{xx} = c_x - de_{1x} - \frac{r_{1s}}{1+s} e_{2x}$$

$$Y_{xy} = c_y - de_{1y} - \frac{r_{1s}}{1+s} e_{2y}$$

$$Y_{xz} = c_z - de_{1z} - \frac{r_{1s}}{1+s} e_{2z}$$

$$(g) \quad Y_{xx} = c_x - de_{1x} + \frac{r_{1s}}{1+s} e_{2x}$$

$$Y_{xy} = c_y - de_{1y} + \frac{r_{1s}}{1+s} e_{2y}$$

$$Y_{xz} = c_z - de_{1z} + \frac{r_{1s}}{1+s} e_{2z}$$

$$(h) \text{ and } (i) \text{ Since } \vec{e}_1 \cdot \vec{e}_2 = \vec{e}_1 \cdot \vec{e}_3 = 0,$$

$$x'_k = - \left[\frac{(r_{kx} - Y_x) e_{2x} + (r_{ky} - Y_y) e_{2y} + (r_{kz} - Y_z) e_{2z}}{(r_{kx} - Y_x) e_{1x} + (r_{ky} - Y_y) e_{1y} + (r_{kz} - Y_z) e_{1z}} \right]$$

$$y'_k = \left[\frac{(r_{kx} - Y_x) e_{3x} + (r_{ky} - Y_y) e_{3y} + (r_{kz} - Y_z) e_{3z}}{(r_{kx} - Y_x) e_{1x} + (r_{ky} - Y_y) e_{1y} + (r_{kz} - Y_z) e_{1z}} \right]$$

END

LITERATURE CITED

1. Harlow, F. H., "The Particle-In-Cell Computing Method for Fluid Dynamics", in Methods in Computational Physics, Eds. Fernbach, and Rotenberg, Eds., Academic Press, New York, 1968, Vol. 2, pp. 319-383.
2. Hader, Charles L., "The Two-Dimensional Hydrodynamic Test Set", Los Alamos Scientific Laboratory report LA-3077 (1956).
3. Hader, Charles L., "The Two-Dimensional Hydrodynamic Test Set, Volume II", Los Alamos Scientific Laboratory report LA-3235 (1956).

DATE FILMED

2 / 17 / 66